

RX ONLY SEMAGLUTIDE SUBLINGUAL SUSPENSION

30 ML VIAL CONCENTRATIONS











STORAGE

Store at controlled room temperature. Beyond-Use-Dating: 90 Days.



DOSING


Begin with a daily sublingual dose of **1 mg for 2-3 weeks**, then gradually increase up to 1.5 mg daily, as needed.



FLAVOR

Hints of smooth caramel-banana provide a pleasant flavor to the suspension.

Request Pricing Catalog Today!



Email: orders@carieboyd.com





PRESCRIBER INFORMATION



RYBELSUS[®]

Our suspension is compounded exclusively with Rybelsus[®], an FDA-approved tablet. This commitment to transparent and responsiblysourced ingredients helps us maintain our devotion to **Compounding with Integrity.**



SUSPENSION -

The base used in our suspension is engineered to handle molecules like semaglutide for sublingual administration. This anhydrous base allows for a **beyond-use date of 90 days** at room temperature.



SUBLINGUAL

The emulsion behavior in the saliva enhances solubility and dispersion. Its permeationenhancing effect works through **mucoadhesion**, increasing contact time with the sublingual mucosa.



DIRECTIONS -

Brush teeth before administration and rinse with warm water, shake the vial well, and administer the dose. Hold under the tongue for **2 minutes**, then swallow. Do not eat or drink for 30 minutes.



INSIGHTS

PCCA consultants have noted **no instances of Gl effects** requiring dose reduction or discontinuation with a daily 1.5 mg sublingual dose. They have also observed consistent weight loss of **one to two pounds** per week.



MONITORING -

Careful evaluation and monitoring of side effects is suggested for dosing beyond 1.5 mg. Monitoring parameters and potential side effects for patients starting semaglutide are the same for any dosage form.

Request Pricing Catalog Today!

Phone: (940) 382-6758

Email: orders@carieboyd.com